

Component Analysis (DHS 10.13)

- Find the right features from given data, in general without any priori information
- Unsupervised approach
- Two popular techniques: Principle Component Analysis and Independent Component Analysis

Principle Component Analysis (DHS 10.13.1)

- A popular technique to project d -dimensional data onto a lower-dimensional subspace in a sum-squared error sense.
- Known also as Karhunen-Loeve Transformation,
- Used in exploratory data analysis
- Uses Eigen Decomposition ($M = V\Sigma V^*$) or Singular Value Decomposition ($M = U\Sigma V^*$).

- Steps

- Compute d -dimensional mean vector μ and $d \times d$ covariance matrix, Σ from data.
- Compute eigenvectors and eigenvalues from the covariance matrix

$$D = V^{-1}\Sigma V$$

where D is the diagonal matrix of eigenvalues of Σ . The matrix V holds eigenvectors that diagonalizes the covariance matrix Σ .

or

Compute eigenvectors and eigenvalues from the characteristic equation (DHS A.2)

Given a $d \times d$ matrix, M , $Mu = \lambda u$

where $u = e_i$ eigenvectors and λ_i associated eigenvalues.

Solve $|M - \lambda I| = \lambda^d + a_1\lambda^{d-1} \dots + a_{d-1}\lambda + a_d = 0$ for each of its d roots λ_j .

For each root solve for eigenvector e_j .

Trace $tr[M] = \sum_{i=1}^d \lambda_i$ and Determinant $|M| = \prod_{i=1}^d \lambda_i$

- ◆ Eigenvector indicate the axis
- ◆ Eigenvalue indicate the amount of variance explained by the axis.
- Sort eigenvectors and eigenvalues in a descending order.
- Choose k eigenvalues and vectors of the large values.
- Generally speaking, with k dimensionality, the signal subspace can be governed. The rest $d - k$ dimensions generally contain noises.
- Form a $d \times k$ matrix A whose column consists of the k eigenvectors.
- Re-represent the data by principal components consists of projecting the data onto the k -dimensional subspace $x' = A^T(x - \mu)$

